


p53 Signaling

CELL CYCLE G2/M DNA DAMAGE SIGNALING PATHWAY







Key Products for p53 Signaling

Visit www.cellsignal.com/p53list for full listing of antibodies, related kits, and reagents.

ANTI	BODIES	APPLICATIONS	REACTIVITY
#4937	53BP1 Antibody	WB, IHC-P, IF-IC	H, Mk
#2675	Phospho-53BP1 (Ser1778) Antibody	WB, IF-IC, F	H, Mk
#2873	ATM (D2E2) Rabbit mAb	WB	H, M
#13050	Phospho-ATM (Ser1981) (D25E5) Rabbit mAb	WB, F	Н
#13934	ATR (E1S3S) Rabbit mAb	WB, IP	H, M, R
#2853	Phospho-ATR (Ser428) Antibody	WB	H, M, R, Mk
	BRCA1 (A8X9F) Rabbit mAb	WB, IP	Н
	Phospho-BRCA1 (Ser1524) Antibody	WB	Н
	cdc2 (POH1) Mouse mAb	WB, IP, IF-IC	H, Mk
	Phospho-cdc2 (Tyr15) (10A11) Rabbit mAb	WB, IP, IF-IC, F	H, M, R, Mk
	Phospho-cdc2 (Thr161) Antibody	WB	H, M, R, Mk
	cdc25A Antibody	WB	H, M
	cdc25B Antibody	WB	H, M, R, Mk
	cdc25C (5H9) Rabbit mAb	WB	H
	Phospho-cdc25C (Ser216) (63F9) Rabbit mAb	WB, IP, IHC-P	H, Mk
	CDK2 (78B2) Rabbit mAb	WB, IP, F	H, M, R, Mk
	Phospho-CDK2 (Thr160) Antibody	WB, IP, F, E-P	H, M, R
	Chk1 (2G1D5) Mouse mAb	WB	H, M, R, Mk
	Phospho-Chk1 (Ser317) (D12H3) XP® Rabbit mAb		H, M, Mk
	Phospho-Chk1 (Ser345) (133D3) Rabbit mAb	WB, IF-IC, F	H, M, R, Mk
	Chk2 (D9C6) XP® Rabbit mAb	WB, IP, IHC-P, IF-IC	Н
	Chk2 (1C12) Mouse mAb	WB, IHC-P, IF-IC	H, Mk
	Phospho-Chk2 (Thr68) (C13C1) Rabbit mAb	WB, IP, IHC-P, F	H
	Phospho-Chk2 (Thr68) Antibody	WB, IP, IF-IC, F	H, Mk
	Cyclin B1 (D5C10) XP® Rabbit mAb	WB, IP, IF-IC, F	H, R
	Cyclin B1 (V152) Mouse mAb	WB, F	H, M, (Hm)
	Cyclin B1 (V152) Mouse mAb (Alexa Fluor® 647 Conjugate)	F	H, M, (Hm)
#4133	Phospho-Cyclin B1 (Ser133) (9E3) Rabbit mAb	WB, IP	Н
	Phospho-Cyclin B1 (Ser147) Antibody	WB	H, M, R, (Pg)
	DNA-PK (3H6) Mouse mAb	WB, IHC-P, IF-IC	H, Mk
	DRAK2 (33D7) Rabbit mAb	WB, IP, IHC-P, F	M
	GADD45 α (D17E8) Rabbit mAb	WB	H, M, R, Mk
	Phospho-MDM2 (Ser166) Antibody	WB	H, M, R
	p21 Waf1/Cip1 (12D1) Rabbit mAb	WB, IP, IHC-P, IF-IC, F	
	p21 Waf1/Cip1 (12D1) Rabbit mAb	IF-IC, F	H, Mk, (Dg)
	(Alexa Fluor® 647 Conjugate)		, , , , , ,
#2527	p53 (7F5) Rabbit mAb	WB, IHC-P, IF-IC, F, ChIP	H, Mk
#2524	p53 (1C12) Mouse mAb	WB, IP, IF-IC, F, ChIP	H, M, R, Hm, Mk
#48818	p53 (D0-7) Mouse mAb	WB, IHC-P, IF-IC, F, ChIP	Н
#9286	Phospho-p53 (Ser15) (16G8) Mouse mAb	WB, IF-IC, F	Н
#8514	Phospho-p53 (Ser15) (16G8) Mouse mAb (PE Conjugate)	F	Н
#9287	Phospho-p53 (Ser20) Antibody	WB	H, Mk
#2521	Phospho-p53 (Ser46) Antibody	WB, IP, IF-IC, F	H, Mk
#9281	Phospho-p53 (Ser392) Antibody	WB	H, M, Mi
#2570	Acetyl-p53 (Lys379) Antibody	WB	H, M
#2525	Acetyl-p53 (Lys382) Antibody	WB	Н
#13109	p63-α (D2K8X) XP® Rabbit mAb	WB, IP, IF-IC, F, ChIP	Н
#9335	Phospho-p90RSK (Ser380) (9D9) Rabbit mAb	WB	H, M, R, Mk, (C)
#11989	Phospho-p90RSK (Ser380) (D3H11) Rabbit mAb	WB, IHC-P, IF-IC	H, M, R, Mk, Mi, (C, X, Z, B, Dg, Pg, Hr)
#11901	WIP1 (D4F7) Rabbit mAb	WB	H, M, R, Mk

RELATED KITS AND REAGENTS


#9947 DNA Damage Antibody Sampler Kit
#7370 PathScan® Total p53 Sandwich ELISA Kit
#7365 PathScan® Phospho-p53 (Ser15) Sandwich ELISA Kit
#7226 PathScan® Acatulated p52 Sandwich ELISA Kit

#12856 PathScan® Stress and Apoptosis Signaling Antibody Array Kit (Chemiluminescent Readout)
#12923 PathScan® Stress and Apoptosis Signaling Antibody Array Kit (Fluorescent Readout)

#8357 Stress and Apoptosis Antibody Sampler Kit


ATM is a serine/threonine kinase that regulates cell cycle checkpoints and DNA repair. Activation of ATM by autophosphorylation on Ser1981 occurs in response to exposed DNA double stranded breaks. ATM kinase regulates a number of proteins involved in cell cycle checkpoint control, apoptosis, and DNA repair.


Phospho-ATM (Ser1981) (D25E5) Rabbit mAb #13050: WB analysis of extracts from HCT 116 cells, untreated (-) or treated with NCS (+), using #13050 (upper) and ATM (D2E2) Rabbit mAb #2873 (lower).



Phospho-p90RSK (Ser380) (D3H11) Rabbit mAb #11989: IF


analysis of NIH/3T3 cells, serumstarved (left), or treated with TPA #4174 (200 nM, 15 min; right), using #11989 (green). Actin filaments were labeled with DY-554 phalloidin (red). Blue pseudocolor = DRAQ5® #4084 (fluorescent DNA dye).






p53 is a tumor suppressor protein which plays a major role in cellular response to DNA damage and other genomic aberrations. DNA damage induces phosphorylation of p53 at Ser15, leading to reduced interaction between p53 and its negative regulator, MDM2. MDM2 regulates p53 accumulation by targeting it for ubiquitination and proteasomal degradation.

Phospho-p53 (Ser15) (1668) Mouse mAb (PE Conjugate) #8514: Flow cytometric analysis of HT-29 cells, untreated (blue) or UV-treated (green), using #8514.



p53 (1C12) Mouse mAb #2524: Chromatin IPs were performed with cross-linked chromatin from 4 x 10° HCT 116 cells treated with UV and either 2.5 µl of #2524 or 2 µl of Normal Rabbit Ig6 #2729 using SimpleChIP® Enzymatic Chromatin IP Kit (Magnetic Beads) #9003 and primers to the designated loci.



Chk2 can phosphorylate p53 at Ser20, which can interfere with p53 and MDM2 binding subsequently preventing ubiquitination and proteasomal degradation of p53.

Chk2 (D9C6) XP® Rabbit mAb #6334: IHC analysis of paraffin-embedded human ovarian serous adenocarcinoma using #6334.



APPLICATIONS & REACTIVITY KEY: WB Western Blotting / IP Immunoprecipitation / IHC Immunohistochemistry / IF Immunofluorescence / F Flow Cytometry / ChIP Chromatin Immunoprecipitation / -IC Immunocytochemistry / -P Paraffin / -F Frozen / E-P Peptide ELISA / H human / M mouse / R rat / Hm hamster / Mk monkey / C chicken / Mi mink / Dm D. melanogaster / X Xenopus / Z zebra fish / B bovine / Dg dog / Pg pig / Sc S. cerevisiae / All all species expected / () 100% sequence homology

© 2016 Cell Signaling Technology, Inc. Cell Signaling Technology, CST, PathScan, SimpleChIP, and XP are trademarks of Cell Signaling Technology, Inc. U.S. Patent No. 5,675,063. All other trademarks are the property of their respective owners.